

---

# Index

---

- AAAtrex, 87, 163, 170, 174  
acetic acid, 73  
acetochlor, 86–87, 169–70, 174  
acifluorfen, 86–87, 174  
ADF (acid detergent fiber), 72  
Agricultural Marketing Service (USDA), 56  
Agricultural Research and Demonstration Centers (University of Illinois), 140  
agroecology, 61  
air-flow fertilizer spreaders, 41–42  
air quality, 62  
alfalfa (*Medicago sativa*)  
  boron deficiency in, 111–12  
  diseases, 75, 199–201  
  for hay/pasture, 60, 65–68, 71–73, 75  
  insect-pest management, 179, 181–83  
  maintenance fertilizer for, 102  
  nutrient levels for, 96, 111  
  temperature thresholds, 2  
alfalfa caterpillars, 182  
alfalfa mosaic, 201  
alfalfa weevils, 179, 181–82  
allelopathy, 49, 59, 61, 161  
alsike clover, 67, 76–77  
ALS (acetolactate synthase) inhibitors, 174–75  
aluminum sulfate, 99  
amines vs. esters, 164–65, 176  
ammonia, 113, 121  
ammonium (NH<sub>4</sub>), 95, 117–18, 121–22, 172  
ammonium acetate test, 94, 104  
ammonium nitrate (NH<sub>4</sub>NO<sub>3</sub>), 121–22  
ammonium sulfate ([NH<sub>4</sub>]<sub>2</sub>SO<sub>4</sub>; AMS), 122, 172  
amylopectin starch, 26  
anhydrous ammonia (NH<sub>3</sub>), 4, 116, 121, 124–26, 128, 134, 136, 139–40  
anions, 95  
anthracnose, 200–202, 204–6  
Aphanomyces root rot, 200–201  
aphids, 39, 41, 179, 191, 193–95, 201, 204  
Apron XL, 199  
armyworms, 179, 181–82, 184, 188–89, 191, 195–96  
aromatic hydrocarbons, 199  
Asian lady beetles, 193  
atmospheric variables. *See* weather and crops  
atrazine, 83, 86–87, 163, 169–70, 173–75  
autotoxicity, 49, 75  
  
*Average Crop, Pasture and Forestry Productivity Ratings for Illinois Soils*, 13  
  
bacteria  
  Bt (*Bacillus thuringiensis*), 17, 184–90  
  coliform, 84  
  nitrification by, 118  
  nodules produced by, 32  
  *Rhizobium*, 76–77  
  *See also* nematodes  
bacterial blight, 200, 204, 207  
bacterial wilt, 179, 200–202  
baled stover/straw, 106  
band seeding, 69  
barium (Ba), 95  
barley, 45, 129–30  
barley yellow dwarf virus. *See* BYDV  
barnyardgrass, 154  
Basagran, 87, 163, 174  
bean leaf beetles, 179, 190–92, 204  
bean pod mottle, 179, 192, 204  
bedding, 145  
bentazon, 87, 163, 174  
Bicep/Bicep II/Bicep Lite/Bicep II Lite/Bicep II Magnum/Bicep Lite II Magnum, 87, 168, 170  
big bluestem, 79–81  
bindweed, 154  
biodiesel, 29  
biofuel crops, 57–58  
bird cherry-oat aphids, 184, 195  
birdseed, 55  
birdsfoot trefoil, 67–68, 76–77  
black chaff, 205, 207  
black cutworms, 179, 188, 190  
blister beetles, 179, 182, 191  
BMPs (best management practices), 87–89  
BMR (brown midrib) trait, 79  
boron (B), 92, 96, 111–12  
botches, 205  
brassicas, 72, 80–81  
Bravo, 199  
Bray P<sub>1</sub> test, 94, 101–2  
broadcast seeding, 41–42, 68–69  
broadleaf weeds, 153  
brown stem rot (BSR), 203–5, 214  
Bt (*Bacillus thuringiensis*) genes, 17, 184–90  
Bt-rootworm hybrids, 50  
buckthorn (*Rhamnus* species), 193  
buckwheat, 56  
*Bulletin*, 180, 182  
bull thistle, 154  
  
bunt, 205–7  
burcucumber, 154  
burdock, 154  
butterweed, 154  
BYDV (barley yellow dwarf virus), 39, 41, 45, 194–95, 205–7  
  
cadmium (Cd), 95  
calcium (Ca), 91, 95, 110  
calcium carbonate (CaCO<sub>3</sub>), 96–99  
Canada thistle, 154  
canola (oilseed rape), 55–56  
canopy cover, 22–23  
carbon dioxide (CO<sub>2</sub>), 9–10, 61–62  
carbon sequestration, 61–62  
caterpillars, 179, 182, 184, 188–89, 191  
CDPs (conservation drainage practices), 147  
CEC (cation exchange capacity), 95, 103–5  
cellulose, 57–58  
cellulosic ethanol, 57  
Cephalosporium stripe, 205–7  
C-FAR (Illinois Council on Food and Agricultural Research), 88  
chemigation, 86  
chickweed, 154  
chloransulam, 86  
chlorine (Cl), 111–12, 167–68  
chlorosis, 112, 211–12, 215  
chromium (Cr), 95  
cicer milkvetch, 77  
Cisne silt loams, 141  
climate. *See* weather and crops  
clopyralid, 86–87, 174  
clover, 60, 67–68, 71, 75–77  
cloverworm, 191  
Cobra, 163, 174  
cocklebur, 140, 154, 158–59, 175  
cold stress, 4  
coliform bacteria, 84  
companion crops, 68, 71, 130  
*Compendium of Research Reports on Use of Non-traditional Materials for Crop Production*, 112  
confectionary sunflower, 55  
conservation tillage  
  crop production with, 137–42  
  crop yields, 140–41  
  deep ripping, 138  
  density/compaction of soil, 138–39  
  erosion/water runoff reduced by, 88, 110, 135  
  fertilization, 93, 107–8  
  
fertilizer considerations in reduced tillage, 139–40  
  machinery and labor costs, 141–42  
  moisture in soil, 137  
  mulch-till, 133, 135, 138, 141  
  no-till, 133–34, 136–42  
  organic matter in soil, 138  
  overview and definition, 134  
  ridge-till, 135–36, 138  
  sampling depth, 93  
  stand establishment, 139  
  strip-till, 109, 134–35, 137, 141  
  temperature of soil, 137, 141  
  weed control, 140  
contamination. *See under* water quality  
continuous cropping, 50  
copper (Cu), 92, 96, 111–12  
  
corn (*Zea mays* L.), 13–26  
  acreage, 49  
  Bt hybrids, 184–90  
  as a C4 crop, 1–2  
  canopy cover, 22–23  
  cobs, 57  
  cold injury, 24  
  cover crops' effects on yield, 59  
  critical time for early growth, 6  
  disease resistance, 17, 201  
  diseases, 179, 201–3, 215–16  
  diversity within a field of, 62–63  
  drought's effects, 24–25  
  ethanol from, 26, 52, 57  
  fertility, 201  
  flex- vs. fixed-ear, 19  
  flooding's effects, 24  
  food-grade, 25–26  
  genetically adjusted, 7, 15–18, 56–57, 184  
  glyphosate-resistant/-tolerant, 50  
  hail's effects, 24  
  heat's effects, 25  
  herbicide injury to, 177  
  high-amylose, 26  
  hybrid GDD requirements, 15, 17  
  hybrid selection, 15–17, 19, 201  
  importance as food crop, 13  
  in-row plant spacing, 22  
  insect-pest management, 179, 183–90  
  insect resistance, 17, 26, 184

- corn (*continued*)
- irrigation, 149, 151
  - key insect pests, 185–90
  - lodging, 17, 185–86
  - maintenance fertilizer for, 102
  - maturity in hybrids, 15–17
  - nematodes in, 215–16, 218
  - nitrogen rates, 113–17
  - nodal root system, 139
  - non-GMO, 26
  - nutriceutical, 26
  - nutrient levels for, 96, 101, 105, 111
  - nutritionally enhanced, 26
  - organic, 26, 56
  - plant development, 13–15
  - plant population, 18–20, 23–24
  - planting date, 17–19, 24, 132
  - planting depth, 18
  - popcorn, 26
  - prices, 20
  - rainfall timing's effects, 6–7
  - replanting, 23–24
  - residue cover from, 136–37
  - row spacing, 19–22
  - seed prices, 20
  - solar energy's effects on
    - carbon fixation, 10
    - vs. sorghum, 45–46
    - vs. soybean, 27–28, 34
  - specialty, 25–26, 56–57
  - stand counting, 23
  - temperature thresholds, 1, 24–25
  - tillage, 140–41, 201
  - waxy, 26
  - weather-related problems, 24–25
  - vs. wheat, 37
  - wheel traffic compaction's effects, 138
  - yield estimates, 25
  - yield of hybrids, 15–16
  - yield trends, 13
  - zinc deficiency in, 112
- corn borers, 17, 179, 183–84, 187–88, 191
- corn earworms, 184, 188–89, 191
- corn flea beetles, 179, 184–85
- corn leaf aphids, 184, 195
- corn rootworms, 17, 26, 50, 52, 131, 179–80, 183–88, 191
- cornstarch, 57
- corrugated roller seeders, 68–69
- cover crops, 58–61, 88–89
- cowpeas, 77
- CP (crude protein), 72
- crabgrass, 154, 158
- cropping systems, 49–63
  - air quality, 62
  - allelopathy, 49, 59, 61, 161
  - alternative crops in Illinois, 54–61
  - biofuels, 57–58
  - buckwheat, 56
  - canola (oilseed rape), 55–56
  - carbon sequestration, 61–62
  - continuous corn, 51–53
  - corn–corn–soybean rotation, 51
  - corn–soybean rotation, 49–53
  - corn–soybean–wheat rotation, 53
  - cover crops, 58–61
  - defined, 49
  - double-cropping, 49, 53
  - and the environment, 61–63
  - intercropping, 49–50
  - limestone rates, 97
  - livestock's role, 50
  - monocropping (monoculture), 50
  - need for change, 63
  - and nitrogen (N) supply, 50–51
  - relay intercropping, 50
  - specialty corn/soybean production, 56–57
  - species diversity, 62–63
  - strip cropping, 50
  - sunflower, 54–55
  - water quality, 62
- crop residue management, 134
- crop yield
  - with conservation tillage, 140–41
  - corn, 13, 15–16, 25, 59
  - and drainage, 144
  - PY (proven-yield) system, 115–16
  - soybean, 13, 36, 59, 140
  - wheat, 13, 37–39, 42–43
- crown rots, 205
- crownvetch, 76–77
- curly dock, 154
- cutworms, 24, 179, 182–84, 188–91, 195
- dandelion, 154
- DDM (digestible dry matter), 72
- Department of Crop Sciences (University of Illinois), 38, 74–75, 214
- dicamba, 87, 163, 174
- 2,4-dichlorophenoxyacetic acid (2,4-D), 87, 163–66, 174, 176
- dicyandiamide (DCD), 123
- dimethenamid, 86–87, 174
- diseases, 197–207
  - in alfalfa, 75, 199–201
  - in corn, 179, 201–3, 215–16
  - corn's resistance, 17, 201
  - fungicides, 197–99, 201, 203, 206
  - and nutrient uptake, 92
  - overview, 197
  - in soybean, 179, 192, 203–5
  - transmitted by insects, 179
  - in wheat, 38–39, 205–7*See also specific diseases*
- ditches, 144–45
- diversity, genetic, 16
- DMI (demethylation inhibitors), 199
- DMI (dry-matter intake), 72
- dormant harvest, 72
- double-cropping, 49, 53
- double-main drainage, 145–46
- downy brome, 154
- downy mildew, 200, 202, 204–5
- DPT (discounted payback time), 146
- drainage
  - benefits, 143–44
  - conservation, 147–49
  - decision, 146–47
  - installation, 147
  - methods, 144–46
  - overview, 143
  - strategy, 147
  - subsurface, 145–46
  - surface, 144–45*Drainage Water Management for the Midwest*, 148
- drought, 8, 24–25
- Drummer Silty clay loam, 146
- drying agents, 73
- Dual/Dual II/Dual II Magnum, 87, 168–70, 174
- ear-attacking caterpillars, 179, 188–89
- Eastern gamagrass, 79–81
- ecological services, 61
- El Niño, 11–12
- emergence scores, 33
- English grain aphids, 195
- environment and cropping systems, 61–63
- EPA (U.S. Environmental Protection Agency), 84, 184
- EPP (early preplant) herbicide applications, 170–71
- ergot, 44, 205
- erosion
  - cover crops for reducing, 58–59
  - factors affecting, 133
  - problems related to, 135–36
  - tillage's effects, 135–36
- esters vs. amines, 164–65, 176
- ethanol, 26, 52, 57–58
- eutrophication, 109
- evapotranspiration, 7–10
- eyespot, 201–2, 205
- Farm Bill, 133
- Feekes scale, 38
- fertigation, 86, 152
- fertilization
  - nitrogen (*see* nitrogen fertilization)
  - with phosphorus (P), 102, 105–7, 110
  - with potassium (K), 103–7, 110
  - in reduced tillage, 139–40
- fertilizers
  - broadcast fertilization, 107–8
  - deep fertilizer placement, 109
  - environmental considerations, 109–10
  - foliar fertilization, 109
  - irrigation systems, 86
  - maintenance, 102–5
  - seed placed (pop-up) fertilization, 108–9
  - site-specific (variable-rate) application, 109
  - soil testing, 86, 92
  - sources, 106–7
  - starter (row) fertilization, 108, 139, 142
  - strip application, 109
  - timing of fertilization, 110
  - 2x2 placement, 108*See also specific fertilizers*
- festulolium, 77–78
- field capacity, 8–9
- Field Crop Scouting Manual*, 180
- field (dry) pea, 54, 77
- first fall frost, 5
- first killing freeze, 16
- “flaxseed.” *See* Hessian flies
- flooding, 24
- flumetsulam, 86–87, 174
- Food Security Act (1985), 133
- foot rots, 205
- forage
  - annual, 78–80
  - quality, 72
  - seeding recommendations, 79
  - species selection, 74–75
  - testing program, 74*See also* grazing; hay and pasture
- Forage Crop Variety Trials in Illinois*, 74–75
- fossil fuels, 61
- foxtail, 154, 158–59, 175, 177
- FRAC (Fungicide Resistance Action Committee), 198–99
- frogeye leaf spot, 203–4
- frost dates, 5
- frost seeding (overseeding), 66, 70
- Fusarium Head Blight Risk Assessment Tool*, 206
- Fusarium head scab*, 38, 205–7
- Fusarium wilt*, 200–201
- garden pea, 77
- genetic diversity, 16
- geology, and groundwater, 85–86
- German (Foxtail) millet, 80
- GIS (geographical information system), 109
- global warming, 61
- glyphosate, 70, 140, 166–68, 173, 175
- GM (genetically modified) traits, 16–18, 26, 29, 56
- Goss's bacterial wilt, 202
- GPS (global positioning system), 94, 109, 211
- grains. *See* barley; oats; rye; sorghum; spelt; triticale; wheat
- grape colapsis, 184, 190–91

- grasses  
 as biofuel crops, 58  
 and conservation tillage, 140  
 as cover crops, 58–59  
 nitrogen rates, 130–31  
 seeding dates, 66  
*See also under* hay and pasture
- grasshoppers, 182, 184, 191, 195  
 grass tetany, 100  
 grazing, 47, 70–71, 74  
 greenbug, 195  
 green cloverworm, 191  
 groundcherry, 154  
 groundwater vulnerability, 85–87  
 growing degree days (GDDs),  
 2–4, 14–15, 17  
 growing degree units (GDUs),  
 2–4  
 grubs, 183–85, 190–93  
 gypsum, 100
- hail, 24  
 hairy vetch, 58–61, 76–77  
 Harness Xtra, 87, 169–70  
 hay and pasture, 65–81  
 annual forages, 78–80  
 cool-season grasses/legumes,  
 65–70, 75–78  
 drying agents/preservatives for  
 hay, 73  
 established stands, 72  
 evaluating older stands, 65–66  
 grass field evaluation, 65  
 hay harvest management,  
 71–73  
 hay-making practices, 73–74  
 hog pastures, 68  
 horse pastures, 68  
 limestone applications, 99  
 nitrogen rates, 130–31  
 pasture-grazing management,  
 74  
 pasture renovation, 70–71  
 seeding recommendations,  
 67–68, 79, 81  
 selecting species, 74–81  
 warm-season grasses, 79–81  
 weed/insect control, 69–70,  
 74 (*see also* insect-pest  
 management)
- Headline, 199, 206  
 heat stress units (HSUs), 4  
 HEL (highly erodible land), 133  
 Helminthosporium diseases,  
 201–2, 205–7  
 hemp dogbane, 140, 154  
 henbit, 154  
 herbicides  
 acid equivalents, 164–67  
 active ingredients, 163  
 additives, 172, 177  
 ammonium fertilizer, 172  
 application rates, 154, 162  
 buffer surfactants/buffer-  
 compatibility agents, 172–73  
 chemical names, 163  
 classified by site of action,  
 173–74  
 common names, 163  
 compatibility agents, 172  
 and conservation tillage, 140  
 crop injury from, 176–77  
 crop oil concentrates (COC),  
 172  
 and drainage, 144  
 drift reduction agents, 172  
 formulation types, 163–70  
 how they work, 173  
 isomers, 167–69  
 nonionic surfactants (NIS),  
 172  
 overview, 162  
 phytotoxicity, 163, 177  
 postemergence, 154–55, 160–  
 62, 171–73, 177  
 premixes, 169–70  
 product labels, 87, 162–63,  
 171–72  
 soil-applied, 157, 170–71  
 stereoisomers, 167–68  
 trade names, 163  
 translocated, 154  
 volatility, 171  
 and water quality, 86–89  
 weed resistance, 173, 175–76
- herringbone drainage, 145–46  
 Hessian flies, 179, 194–96  
 Hessian fly-free dates, 39–41,  
 43, 195  
 HFS (highly fermentable starch),  
 26  
 hidden-hunger checkups, 94–95  
*Hoplolaimus galeatus*, 216  
 hopperburn, 182–83  
 horsetail, 154  
 horseweed, 154, 175  
*How a Corn Plant Develops*, 13  
 hydrogen (H), 95, 113, 121, 166  
 hypomagnesaemia, 100
- ICP (inductively coupled plasma)  
 emission spectroscopy, 94, 101  
 Illinois Department of  
 Agriculture, 97, 141  
 Illinois Department of Public  
 Health, 84–85  
 Illinois Department of  
 Transportation, 97  
*Illinois Drainage Guide (Online)*,  
 146, 148  
 Illinois EPA (Environmental  
 Protection Agency), 83, 86  
*Illinois Integrated Water Quality  
 Report and Section 303d  
 List—2008*, 83  
*Illinois Pesticide Applicator  
 Training Manual 39-2: Field  
 Crops*, 179  
 Illinois State Geological Survey,  
 151  
 Illinois State Water Survey, 85,  
 151
- Illinois Voluntary Limestone  
 Program Producer  
 Information*, 97  
 Illinois Well Construction Code,  
 85  
 imazethapyr, 87, 173–74  
 Indiangrass, 79–81  
 inoculation of legumes, 32, 70,  
 76–77  
 insecticides  
 for alfalfa, 182  
 chloronicotinyl, 184–85, 188,  
 190, 194–96  
 foliar-applied, 192–96  
 seed-applied, 41, 184–85, 188,  
 190, 194–96  
 soil-applied, 185
- insect-pest management, 179–96  
 curative, 180  
 developing strategies, 180–81  
 for hay and pasture, 69–70, 74  
 IPM (integrated pest  
 management), 88, 180, 183,  
 187  
 overview, 179–80  
 preventive, 180  
*See also* pesticides and specific  
 insects and other pests
- interceptor drains, 144  
 intercropping, 49–50  
 interseeding, 70–71  
 IPM (integrated pest  
 management), 88, 180, 183,  
 187  
 IRM (insect resistance  
 management) strategies, 184,  
 188  
*See also* insect-pest  
 management
- iron (Fe), 92, 95–96, 99, 107,  
 111–12  
 iron sulfate, 99  
 IRR (internal rate of return), 146  
 irrigation  
 benefits, 148–51  
 decision, 151–52  
 fertigation, 86, 152  
 management requirements,  
 152  
 scheduling, 152  
 subsurface, 151–52
- Japanese beetles, 179, 184,  
 191–93  
 Japanese millet, 80  
 Jerusalem artichoke, 154  
 jimsonweed, 140, 154  
 johnsongrass, 154
- kale, 72, 80–81  
 karst (limestone region), 86  
 Kentucky bluegrass, 68, 77–78,  
 130–31  
 kochia, 154, 157, 175  
 kura clover, 76–77
- lactofen, 163, 174  
 ladino clover, 60, 75  
 LAI (acres of leaves per acre of  
 crop), 28  
 lambsquarters, 154, 158, 173, 175  
 La Niña, 11  
 last spring frost, 5  
 late-summer seedings, 66  
 leaf blight, 179, 201–2  
 leafhoppers, 71, 75–76, 179,  
 181–83, 191  
 leaf spot, 39, 77, 198, 200–204  
 legumes, 32  
 as cover crops, 59–60  
 for hay and pasture, 65–70,  
 75–78  
 inoculation, 32, 70, 76–77  
 seeding dates, 66  
 species, 75–77  
*See also specific legumes, such  
 as soybean*
- lespedeza, 66–67, 76–77, 96–97  
 lime, 94, 99  
 limestone  
 applying, 99  
 calcitic (CaCO<sub>3</sub>), 100, 111  
 dolomitic (CaMg[CO<sub>3</sub>]<sub>2</sub>), 100,  
 111  
 ENV (effective neutralizing  
 value), 96–99  
 low-Mg, 100  
 for pH neutralizing, 96–99  
 rate calculation, 97–99  
 rate recommendations, 97  
 sampling, 97  
 little barley, 154  
 lupines, 77
- magnesium (Mg), 91, 95, 111  
 manganese (Mn), 92, 95–96, 99,  
 111–12  
 manure, 115, 125, 131–32  
 Maxim, 199  
 MBC (methyl benzimidazole  
 carbamates), 199  
 MCLs (maximum contaminant  
 levels), 83  
 meadow fescue, 77–78  
 mefenoxam, 203–4  
 Mehlich-3 test, 94, 101, 104  
 mercury (Hg), 95  
 metalaxyl, 203–4  
 metolachlor, 87, 168–70, 174  
 metribuzin, 86–87, 174  
 microbes, 50, 59, 117, 121  
 milk fever, 100  
 milkweed, 154  
 milo. *See* sorghum  
 mineralization, 106, 117–18, 132  
 miscanthus (*Miscanthus x  
 giganteus*), 2, 58  
 moldboard plowing, 138  
 molybdenum (Mo), 96, 111–12,  
 128

- monoammonium phosphate (MAP) vs. diammonium phosphate (DAP), 106, 110, 123
- monocropping (monoculture), 50
- morningglory, 154, 158
- multiple cropping. *See* double-cropping
- muriate of potash (potassium chloride; KCl), 107
- musk thistle, 154
- narrow-row planters, 142
- National Agricultural Statistics Service, 20
- National Corn Growers Association, 184
- National Forage Testing Association, 72
- National Weather Service, 10–11
- Natural Resources Conservation Service (NRCS), 83, 134
- Natural Resources Conservation Service Field Office Technical Guide*, 133
- NBPT (thiophosphoric triamide), 124
- NDF (neutral detergent fiber), 72–73
- NDVI (normalized difference vegetation index), 120–21
- nematodes, 209–18
  - corn-parasitic, 215–16, 218
  - dagger, 216–18
  - damage thresholds, 217–18
  - how they damage plants, 209–12
  - lance, 216–18
  - lesion, 215–16, 218
  - needle, 216–18
  - overview, 209
  - pin, 215, 218
  - root-knot, 215, 218
  - SCN (soybean cyst nematodes), 55, 203–4, 209–15, 218
  - scouting for, 211–12
  - sting, 215, 218
  - stun, 215
- Nematology Lab (University of Illinois), 214–15
- nightshade, 140, 158–59, 175
- La Niña, 11–12
- El Niño, 11–12
- nitrapyrin (N-Serve), 88, 123
- nitrate (NO<sub>3</sub>)
  - contamination, 83–84
  - early spring nitrate nitrogen test, 119
  - as fertilizer, 121
  - leaching, 88, 118–19
  - PSNT (pre-sidedress nitrate test), 119–20
  - uptake, 117
- nitrite (NO<sub>2</sub>), 118
- nitrogen (N), 113–32
  - availability, estimating, 119–21
  - availability, factors affecting, 117–19
  - barley, rates for, 129–30
  - carryover, and idled acres, 131
  - corn rates, 113–17
  - crop color sensing technology, 120–21
  - cycle of, 117
  - denitrification, 8–9, 118
  - early spring nitrate nitrogen test, 119
  - fertigation, 152
  - fixation, 32
  - grass hay, rates for, 130–31
  - immobilization, 117
  - ISNT (Illinois soil nitrogen test; amino sugar-N test), 119
  - leaching, 8–9, 86, 118–19
  - in manure, 115, 131–32
  - mineralization, 117–18
  - MRTN (maximum return to N), 114–16, 124
  - nitrification, 4, 118
  - nitrification inhibitors, 123–24, 130
  - oats, rates for, 129–30
  - overview, 113
  - pasture, rates for, 131
  - and planting date, 132
  - plant tissue testing, 120
  - previous crop, adjustments for, 131–32
  - PSNT (pre-sidedress nitrate test), 119–20
  - PY (proven-yield) system, 115–16
  - rate adjustments, 131–32
  - RTN (return to N) values, 114–15
  - soybean, rates for, 128–29
  - SPAD readings, 120–21
  - total soil nitrogen test, 119
  - wheat, rates for, 129–30
- nitrogen fertilization, 121–28
  - amendments, 123–24
  - ammoniated phosphate, 123
  - ammonium nitrate (NH<sub>4</sub>NO<sub>3</sub>), 121–22
  - ammonium sulfate ([NH<sub>4</sub>]<sub>2</sub>SO<sub>4</sub>), 122
  - anhydrous ammonia (NH<sub>3</sub>), 4, 116, 121, 124–26, 128, 134, 136, 139–40
  - application methods, 126–28
  - application timing, 124–26
  - nitrogen solutions, 122
  - organic-N, 123
  - urea (CO[NH<sub>2</sub>]<sub>2</sub>), 121–22, 124, 128
- nitrous oxide (NO<sub>2</sub>), 118
- nodules, nitrogen fixation in, 31
- no-till drills, 69–71, 136
- NPV (net present value), 146
- nutriceuticals, 26
- nutrients. *See* soil pH and crop nutrients *and* specific nutrients
- Nutrient Standards Forum (University of Illinois–Springfield, 2007), 88
- nutsedge, 154, 159
- oats
  - broadcast vs. drilled, 45
  - as a companion crop, 68, 130
  - disking the stalks, 45
  - for hay and pasture, 77
  - herbicide sensitivity, 45
  - maintenance fertilizer for, 102
  - nitrogen rates, 129–30
  - nutrient levels for, 101
  - planting dates, 45
  - seed treatment with fungicides, 45
  - spring variety, 44–45
  - test weight, 44
  - varieties, 45
  - winter variety, 45
- oilseed sunflower, 54–55
- Olsen (sodium bicarbonate) test, 94, 101
- Optimum Crop Productivity Ratings for Illinois Soils*, 13
- orchardgrass, 66–68, 77, 130–31
- organic-N fertilizers, 123
- ozone, 62
- P. brachyurus*, 216
- P. hexincisus*, 216
- P. zea*, 216
- Pacific Ocean, 11
- panicum, 158
- parallel drainage, 145–46
- paraquat, 70–71, 174
- parasites. *See* nematodes
- pasture. *See* grazing; hay and pasture
- PCNB, 199
- PEAQ (“Predictive Equations for Alfalfa Quality”), 72–73
- pearl millet, 79
- Pedersen, Palle, 29
- pennycress, 154
- Performance of Commercial Corn Hybrids in Illinois*, 17
- Performance of Commercial Soybeans in Illinois*, 28
- pesticides, 83, 85, 88, 163
  - See also* insecticides
- Pest Management of Alfalfa Insects in the Upper Midwest*, 182–83
- pests
  - in corn, key, 185–90
  - defoliators, 191–93, 195
  - key pests, 179
  - and nutrient uptake, 92
  - population density, 180–81
  - predators, 180
  - soybean pest list, 191
  - See also* insect-pest management; nematodes; *and* specific pests
- pH. *See* soil pH and crop nutrients
- phenylamides, 199
- phenylpyrroles, 199
- phosphorus (P)
  - fertilization, 102, 105–7, 110
  - in manure, 132
  - for pasture renovation, 70
  - and pH, 99
  - as a pollutant, 109
  - recommendations, 101–2
  - removal, 101, 106
  - soil testing, 93–95, 101
  - solubility, 106–7
  - stratification, 139
  - supplying power, 100–101, 105
- phosphorus pentoxide (P<sub>2</sub>O<sub>5</sub>), 102, 105–7, 109
- photoperiod sensitivity, 27
- photosynthesis, 9–10, 17, 24–25, 61–62, 71, 92, 173–74
- physical climate forecasting, 11
- Phytophthora root rot, 200–201, 203–4
- PI (profitability index), 146
- pigweed, 154, 159, 175
- Pioneer Hi-Bred International, 29
- Planning Your Well: Guidelines for Safe, Dependable Drinking Water*, 84
- plant analysis, 95
- Plant Clinic (University of Illinois), 197, 215
- PLS (pure live seed), 67
- POC (petroleum oil concentrate), 172
- pod blight, 203, 205
- poison hemlock, 154
- pokeweed, 154
- pollution, 62
- popcorn, 26
- potash, white vs. red, 107
- potassium (K)
  - as a cation, 95
  - as a drying agent, 73
  - fertilization, 103–7, 110
  - in manure, 132
  - overview, 103
  - for pasture renovation, 70
  - recommendations, 103–5
  - removal, 106
  - soil testing, 93–95, 103
  - stratification, 139
- potassium oxide (K<sub>2</sub>O), 102, 105–6, 108–9
- potato leafhoppers, 75–76, 179, 181–83, 191
- powdery mildew, 204–7
- PPI (preplant incorporated) herbicide applications, 170
- PPO (protoporphyrinogen oxidase) inhibitors, 174–75
- precipitation, 5–8, 119, 150

- PRE (preemergence) herbicide applications, 170
- preservatives, 73
- prickly lettuce, 154
- propionic acid, 73
- Prosaro, 199
- Prowl, 163, 174
- prussic acid (hydrocyanic acid, or HCN), 47, 78–79
- purple deadnettle, 154
- purple seed stain, 203, 205
- Pursuit, 166, 173–74
- Pythium, 201, 205
- QoI (quinone outside inhibitors), 199
- quackgrass, 154
- Quadris, 206
- Quilt, 206
- ragweed, 154, 158–60, 175
- rain, 5–7, 119  
*See also* precipitation
- random drainage, 145
- rape, 55, 68, 80–81
- rapeseed, 55
- Raptor, 163, 174
- recycled paper, 58
- red clover, 60, 66–68, 71, 75
- redtop, 67
- reed canarygrass, 67, 77, 130–31
- relative maturity (RM), 16–17
- relay intercropping, 43, 50
- RFQ (relative forage quality), 72
- RFV (relative feed value), 72–73
- Rhizobium* bacteria, 76–77
- riparian doctrine, 151
- root development and drainage, 144
- root rots, 200–201, 203–4
- Roundup, 173–74  
*See also* glyphosate
- roundworms. *See* nematodes
- runoff water, 59, 62, 83, 86, 88–89, 109–10, 123, 144
- rusts, 200, 202–7
- rutabaga, 80–81
- rye  
 Balbo, 44  
 as a cover crop, 58–60  
 ergot fungus in, 44  
 Hancock, 44  
 planting date, 44  
 Rymin, 44  
 Spooner, 44  
 spring vs. winter varieties, 44  
 winter-/drought-hardiness, 44
- ryegrass, 58, 66–68, 77–78
- Safe Drinking Water: Testing and Treating Home Drinking Water*, 84
- Safe Drinking Water hotline, 84
- sainfoin, 77
- Salvo, 163
- saturated fats, 36
- Sclerotinia stem rot (white mold), 56, 200, 204–5
- SCN (soybean cyst nematodes), 55, 203–4, 209–15, 218
- SDS (sudden death syndrome), 204–5, 214
- Sealing an Abandoned Well*, 85
- sedimentation, 136
- seedcorn maggots, 184, 190
- seed drop calculator, 31, 40
- seeding no-till, 69
- Septoria diseases, 39, 203–7
- sequential cropping. *See* double-cropping
- shattercane, 154, 175
- shepherd's-purse, 154
- shoot-to-root ratio, 6
- sidedressing, 120, 123, 125–28, 132, 152
- sidewall smearing, 139
- silt loam/silty clay loam, 9, 141, 146
- simazine, 86–87, 173–75
- sinkholes, 86
- small grains. *See* barley; oats; rye; sorghum; spelt; triticale; wheat
- smartweed, 154, 158–59
- smooth bromegrass, 67–68, 77, 130–31
- smut, 201–2, 205–7
- snow, 5–6
- sodium (Na), 95, 107, 163
- sodium carbonates, 73
- soil  
 aeration, 143  
 characteristics, and groundwater, 85  
 compaction, 8, 18, 91, 99, 110, 124–25, 127, 138–40  
 conservation compliance, 133–34  
 dry, 9  
 field, 92–93  
 management overview, 133  
 moisture, 8–9, 91  
 runoff, 62  
 stable organic matter in, 61–62  
 structure, 144  
 temperature, 4, 91–92, 125, 143  
 wet, 8–9  
*See also* conservation tillage; drainage; dry soil; soil moisture; soil pH and crop nutrients; tillage; wet soil
- Soil and Water Conservation Districts, 151
- soilborne mosaic, 39, 205
- soil pH and crop nutrients, 91–112
- baled stover/straw, 106
- calcium–magnesium balance, 100
- CEC (cation exchange capacity), 95, 103–5
- environmental considerations, 109–10
- micronutrients, 94–95, 111–12
- nontraditional products, 112
- nutrient availability, estimating, 92–95
- nutrient availability, factors affecting, 91–92
- nutrient availability, relative to pH, 96
- overview, 91
- pH effects on plant growth, 95–97
- pH goals, 96
- pH lowering (acidifying), 99–100
- phosphorus (P), 100–102, 105–6
- pH raising (liming), 96–97, 139 (*see also* limestone)
- pH rate calculation, 97–99
- plant analysis, 95
- potassium (K), 103–6
- secondary nutrients, 94–95, 110–11
- soil testing, 92–96, 100–106, 108–9  
*See also* fertilizers; and specific nutrients
- Soils of Illinois*, 13
- SOILS Project, 141
- solar radiation, 8–10
- sorghum (*Sorghum bicolor*; grain sorghum)  
 vs. corn, 45–46  
 fertilization, 46  
 forage, 78–79  
 grazing, 47  
 harvesting/storage, 47  
 hybrid selection, 46  
 maintenance fertilizer for, 102  
 marketing, 47  
 plant population, 46  
 planting date, 46  
 planting depth, 46  
 row spacing, 46  
 temperature thresholds, 2  
 weed control, 47
- sorghum, sweet, 46, 79
- sowthistle, 154
- soybean (*Glycine max*, Merrill), 27–36  
 acreage, 49  
 canopy cover, 28, 30, 33  
 vs. corn, 27–28, 34  
 cover crops' effects on yield, 59  
 defoliation, 191–93  
 diseases, 179, 192, 203–5  
 double-cropped, 30, 34 (*see also under* cropping systems)  
 fertilizer's effects on nitrogen fixation, 32  
 glyphosate-resistant/-tolerant, 34, 50, 134  
 GMO (genetically modified) vs. non-GMO, 29–30, 35, 56–57  
 high-oleic, 36  
 high-saturated-fat, 36  
 high-yield, 36  
 inoculation, 32  
 insect-pest management, 179, 190–94  
 iron deficiency in, 112  
 low-linolenic, 36  
 low-saturated-fat, 36  
 maintenance fertilizer for, 102  
 manganese deficiency in, 112  
 MGs (maturity groups), 28  
 natto-type, 36  
 nitrogen rates, 128–29  
 nutrient levels for, 96, 101, 105, 111  
 oil from, 29, 36  
 organic, 36, 56  
 pest list, 191  
 plant development, 27–28, 34  
 plant population, 30  
 planting date, 29–30  
 planting depth, 33  
 planting rate and seed issues, 30–31  
 protein from, 29  
 rainfall timing's effects, 6–7  
 relay intercropping, 43  
 replanting, 34–35  
 residue cover from, 137  
 Roundup Ready, 29  
 row width, 31, 33, 35, 142, 205  
 SCN (soybean cyst nematodes) in, 55, 203–4, 209–15, 218  
 seed quality/testing, 31–32  
 seed size, 32  
 solar energy's effects on carbon fixation, 10  
 specialty, 35–36, 56–57  
 temperature thresholds, 2  
 thin-line vs. bush-type, 33  
 tillage's effects on yield, 140  
 tofu-type (clear hilum), 36  
 variety selection, 28–29  
 wheel traffic compaction's effects, 138  
 yield trends, 13
- soybean aphids, 190–91, 193–94, 204
- Soybean Growth and Development*, 27
- Soybean Rust IPM PIPE, 203
- SPAD meter readings, 120–21
- species diversity, 62–63
- spelt, 44
- spider mites, 179, 191, 193–94
- spinner-type seeders, 66
- split-row planters, 142
- spring seedings, 66, 71–72
- sprinklers, overhead, 152
- SRI (strategic research initiative), 88
- Stagonospora diseases, 205–7

- stalk borers, 184, 191, 195  
 statistical climate forecasting, 11  
 stay-green effect, 199  
 stem blight, 203, 205  
 Stewart's bacterial wilt, 179, 202  
 Stratego, 206  
 strip cropping, 50  
 stunting, 190, 193, 210–12, 215–16  
 sudangrass, 78–79  
 sugarcane, 57  
 sulfentrazone, 86–87, 174  
 sulfur (S), 92, 96, 99–100, 110–11, 122  
 sunflower, 54–55, 158  
 sunshine, 9  
   *See also* solar radiation  
 surface-water contamination, 86–89  
 sweep nets, 183  
 sweetclover, 76–77  
 sweet sorghum, 46, 79  
 switchgrass, 47, 58, 79–81
- take-all, 205–7  
 tall fescue, 67–68, 77–78, 130–31  
 tan spot, 205, 207  
 teasel, 154  
 teff, 79–80  
 temperature, 1–5  
   forecasting, 11  
   growing degree units (GDUs), 2–4  
   sea-surface, 12  
   of soil, 4, 91–92, 125, 143  
   stress, 4  
   thresholds, 1–2  
 thistle caterpillars, 191  
 tile drainage, 62  
 tillage, 133–42  
   clean, 135  
   conventional, 135, 137  
   overview, 133  
   residue cover affected by, 136–37  
   soil erosion affected by, 135–36  
   trends, 134  
   of wheat, and previous crop, 41  
   *See also* conservation tillage  
 Timothy, 67–68, 77  
 tipburn, 182–83  
 TMDLs (total maximum daily loads), 86–87  
 Topsis M, 199  
 trans fats, 36, 57  
 triazines, 174–75  
 triticale, 44  
 trumpetcreeper, 154  
 turnips, 72, 80–81  
 twospotted spider mites, 179, 191, 193–94  
 UAN (urea-ammonium nitrate), 122, 124, 127–28, 130, 172
- UPT (undiscounted payback time), 146  
 urea (CO[NH<sub>2</sub>]<sub>2</sub>), 121–22, 124, 128, 142  
 urease inhibitors, 124, 140  
 USDA National Soybean Germplasm Collection (University of Illinois–Urbana), 214  
 USDA rules regarding organic crops, 56  
 USEPA (U.S. Environmental Protection Agency), 83, 87–88  
*Utilizing Herbicide Site of Action to Combat Weed Resistance to Herbicides*, 173
- Valor, 163, 174  
 velvetleaf, 140, 154, 158–59  
 vernalization, 40  
 verticillium wilt, 200–201  
 Vitavax, 199  
 V (vegetative-stage) numbers, 27  
 VOC (vegetable oil concentrate), 172  
 VRT (variable-rate technology), 109
- waterhemp, 140, 154, 157–59, 175  
 water management, 143  
   *See also* drainage; irrigation  
 water pH test, 94  
 water quality  
   BMPs (best management practices), 87–89  
   boiling of water, 84  
   chemical properties/selection, 86  
   consumer confidence reports, 83–84  
   in cropping systems, 62  
   drinking-water standards, 83  
   groundwater vulnerability, 85–87  
   irrigators, 86  
   nutrient standards, 87–88  
   overview, 83  
   point-source contamination, 84–85  
   sealing abandoned wells, 85  
   surface-water contamination, 86–89  
   testing private wells, 84  
 water table level, 147, 149  
 Water Use Act (1983), 151  
 weather and crops, 1–12  
   El Niño/La Niña's effects on Illinois climate, 11–12  
   evapotranspiration, 7–10  
   forecasting, 10–12  
   growing degree units (GDUs), 2–4  
   growing season length, 5  
   normal weather conditions, defined, 1
- overview, 1  
 precipitation, 5–7  
 soil moisture, 8–9  
 soil temperature, 4  
 solar radiation, 8–10  
 temperature, 1–5  
 weed management, 153–77  
   annuals, 153–55  
   biennials, 153–54, 156  
   chemical (*see* herbicides)  
   and conservation tillage, 140  
   cropping and cultural practices, 159, 162  
   emergence sequences, 158  
   hay and pasture, 69–70, 74  
   interference, 157–62  
   lessons from research, 160–62  
   mechanical practices, 162  
   overview, 153  
   perennials, 154, 156  
   scouting fields for weeds, 154, 158  
   for sorghum, 47  
 weed variables, 159–60  
*Weeds of the Great Plains*, 153  
*Weeds of the North Central States*, 153  
*Weeds of the Northeast*, 153  
 wells  
   abandoned, 85  
   depths, and groundwater, 86  
   notification of drilling, 151  
   testing, 84  
 wheat (*Triticum aestivum*), 37–47  
   acreage, 49  
   vs. corn, 37  
   as a cover crop, 58–59  
   in the cropping system, 37 (*see also under* cropping systems)  
   diseases, 38–39, 205–7  
   diseases in, 39  
   double-cropped, 34, 37, 42–43  
   erosion control by, 37–38  
   fertility, 206  
   hard vs. soft, 37–38  
   insect-pest management, 179, 194–96  
   insects in, 39, 41  
   intensive management in Illinois vs. Europe, 42  
   maintenance fertilizer for, 102  
   management for best yields in Illinois, 42  
   marketing classes of, 37  
   nitrogen rates, 129–30  
   nutrient levels for, 96, 101  
   plant development, 38  
   planting date, 39–40, 43, 206  
   planting depth, 41–42  
   relay intercropping, 43  
   row spacing, 42  
   seeding rate, 40–41  
   seed treatment with fungicides/insecticides, 41, 194–95, 206  
   spring variety, 43–44  
   straw yield from, 38–39, 43
- temperature thresholds, 2  
 tillage, 41, 140, 205–6  
 variety selection, 38–39  
 volunteer, 41  
 white vs. red, 37–38  
 winter-hardiness, 39–40, 42  
 winter variety, 37–38, 40  
 yield trends, 13, 37  
 wheat curl mites, 179, 195  
 wheat streak mosaic, 39, 179, 195, 202, 205–7  
 white clover, 66–68, 75–76  
 white grubs, 183–85, 190–91  
 wild carrot, 154  
 wild proso millet, 158  
 wilting point, 9  
*Winter Survival, Fall Dormancy and Pest Resistance Ratings for Alfalfa Varieties*, 75  
 wirestem muhly, 154  
 wireworms, 183–85, 190–91  
 wood processing wastes, 58  
 woolly cupgrass, 158  
 woollybear caterpillars, 184, 191  
 World Meteorological Organization, 1
- yellow rocket, 154  
 Yukon, 87, 163
- zinc (Zn), 92, 95–96, 99, 111–12